arXiv:math/0403535v1 [math.AC] 31 Mar 2004

MONOMIAL IDEALS ARISING FROM DISTRIBUTIVE LATTICES

XINXIAN ZHENG

ABSTRACT. The free resolution and the Alexander dual of squarefree monomial ideals associated with certain subsets of distributive lattices are studied.

INTRODUCTION

Let $\mathscr{L}$ be a finite distributive lattice. By Birkhoff's fundamental structure theorem, there is a unique poset (partially ordered set) P such that $\mathscr{L}$ is isomorphic to the poset $\mathscr{J}(P)$ consisting of all poset ideals (including the empty set) of P, ordered by inclusion. In fact, P can be chosen as the set of all join-irreducible elements of $\mathscr{L}$. Let K be a field and $S = K[\{x_p, y_p\}_{p\in P}]$ the polynomial ring in $2|P|$ variables over K with $\deg x_p = 1$ and $\deg y_p = 1$ for all $p \in P$, and let $\mathscr{S} \subset \mathscr{L}$ be any subset of $\mathscr{L}$. The Hibi ideal $H_{\mathscr{S}}$ associated with $\mathscr{S}$ is the monomial ideal in S generated by the monomials u_p with $p \in \mathscr{S}$, where $u_p = x_{\ell(p)} y_{P\setminus \ell(p)}$ and where $\ell(p)$ is the principal poset ideal $\{q \in P\colon q \leq p\}$ in P.

In [8] it is shown that for any poset ideal $\mathscr{I}$ of $\mathscr{L}$, the Hibi ideal $H_{\mathscr{I}}$ has a linear resolution. In this article, we consider more generally the ideal $H_{\mathscr{S}}$ where $\mathscr{S}$ is a segment of $\mathscr{L}$ (see Definition 2.1). For example, any poset ideal $\mathscr{I}$, or any poset coideal $\mathscr{J}$ of $\mathscr{L}$, as well as their intersection are segments in $\mathscr{L}$. In the third section we describe in Theorem 3.8 when $H_{\mathscr{I}} \cap H_{\mathscr{J}} = H_{\mathscr{I}\cap\mathscr{J}}$, and in Theorem 3.9 it is said when this ideal has a linear resolution. In particular this answers a question which was raised in [8], see Corollary 3.15 and 3.16. We also show in Theorem 3.10 that the ideal $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has always a linear resolution, if $\mathscr{I} \cup \mathscr{J} = \mathscr{L}$ and $\mathscr{I} \cap \mathscr{J} = \emptyset$.

Let G be a Cohen–Macaulay bipartite graph on the vertex set $V \cup V'$ with $V \cap V' = \emptyset$ and $|V| = |V'| = n$, and $S = K[x_1, \ldots, x_n, y_1, \ldots, y_n]$ the polynomial ring over a field K. In [6, Theorem 2.4] the authors showed that the vertices $V = \{x_1, \ldots, x_n\}$ and $V' = \{y_1, \ldots, y_n\}$ can be labeled such that there exists a partial order $<$ on V with the property that $\{x_i, y_j\}$ is an edge of G if and only if $x_i \leq x_j$. Moreover it is shown that for $P = (V, <)$ the distributive lattice $\mathscr{J}(P)$ satisfies $H^*_{\mathscr{J}(P)} = I(G)$. Here, for any subset $\mathscr{S}$ of $\mathscr{J}(P)$ we denote by $H^*_{\mathscr{S}}$ the defining ideal of the Stanley–Reisner of the Alexander dual of Γ, where Γ is the simplicial complex defined by the equation $H_{\mathscr{S}} = I_\Gamma$.

Later, in [8] the authors considered more generally simplicial complexes Δ on the vertex set $V \cup V'$ with $V \cap V' = \emptyset$ and $|V| = |V'|$ such that

(1) there is no $F \in \mathscr{F}(\Delta)$ with $F \subset V$,
(2) $G = \{F \in \mathscr{F}(\Delta)\colon F \cap V \neq \emptyset, \quad F \cap V' \neq \emptyset\}$ is a Cohen–Macaulay bipartite graph with no isolated vertex,

---

1991 *Mathematics Subject Classification.* 16D25, 16E05, 06D50, 06D99.

and studied when the facet ideal $I(\Delta)$ of Δ is Cohen-Macaulay. As a further generalization we consider in the second section of this article simplicial complexes Δ satisfying only condition (2), and show (Theorem 2.4) that Δ is unmixed and each minimal vertex cover of Δ has cardinality n if and only if there exists a segment $\mathscr{S}$ of some distributive lattice $\mathscr{L}$ such that $H^*_{\mathscr{S}} = I(\Delta)$.

I would like to thank Jürgen Herzog for many helpful comments and discussions.

1. PREPARATIONS

In this section we recall some basic facts on lattices and simplicial complexes and fix some notation. As a general reference for posets and lattices we refer the reader to [5] and [9], and to [10], [2] and [11] concerning simplicial complexes, Stanley-Reisner and facet ideals.

Let P be any finite poset (partially ordered set), and let $\alpha, \beta \in P$ with $\alpha \leq \beta$. The set

$$[\alpha,\beta] = \{\gamma \in P : \alpha \leq \gamma \leq \beta\}$$

is called the *interval* between α and β in P.

Let P be a poset and $\alpha, \beta \in P$. If $\alpha < \beta$ and for each element $\gamma \in P$ with $\alpha \leq \gamma \leq \beta$, we have either $\gamma = \alpha$ or $\gamma = \beta$, then we say β *covers* α, or α is a *lower neighbor* of β, or β is an *upper neighbor* of α.

An element in a poset P may have more than one upper neighbor (resp. lower neighbor) or have no upper neighbor (resp. lower neighbor). An element in a poset P which has exactly one lower neighbor is called a *join irreducible element* of P. The set of all join irreducible elements with the induced order is a poset, called the *join irreducible subposet* of P. Conversely, an element in a poset P which has exactly one upper neighbor is called a *meet irreducible element* of P.

A *chain* is a poset in which any two elements are comparable. A subset C of a poset P is called a *chain* if C is a chain when regarded as a subposet of P. The *length* $\ell(C)$ of a finite chain is defined by $\ell(C) = |C| - 1$. The *length* (or *rank*) of a finite poset P is $\ell(P) := \max\{\ell(C) : C \text{ is a chain of } P\}$. If every maximal chain of P has the same length r, then we say P is *graded of rank* r. In this case there is a unique *rank function* $\rho : P \to \{0, \ldots, r\}$ such that $\rho(\alpha) = 0$ if α is a minimal element of P, and $\rho(\beta) = \rho(\alpha) + 1$ if β covers α in P. If $\rho(\alpha) = i$, then we say α has *rank* i.

Later, we need the *dual poset* of P. This is the poset $\widetilde{P}$ on the same set as P, but such that $\alpha \leq \beta$ in $\widetilde{P}$ if and only if $\beta \leq \alpha$ in P.

A *lattice* is a poset $\mathscr{L}$ for which each pair of elements α and β has a least upper bound (called the *join* of α and β, denoted by $\alpha \vee \beta$) and a greatest lower bound (called the *meet* of α and β, denoted by $\alpha \wedge \beta$).

One sees immediately from the definition that in a lattice $\mathscr{L}$, there is a unique element μ satisfies that $\mu \geq \alpha$ for any $\alpha \in \mathscr{L}$. This element is called the *maximum* of $\mathscr{L}$, and denoted by $\hat{1}$. Similarly, there is a unique element ν satisfies $\nu \leq \alpha$ for any $\alpha \in \mathscr{L}$. This element is called the *minimum* of $\mathscr{L}$, and denoted by $\hat{0}$.

A *poset ideal (coideal)* of a poset P is a subset I of P such that if $\alpha \in I$ and $\beta < \alpha$ ($\beta > \alpha$), then $\beta \in I$. The maximal (minimal) elements in I are called the *generators* of I. The set of generators is denoted by $G(I)$.

Remark 1.1. Let $I \subset P$. Then the following conditions are equivalent:

1. I is a poset ideal (coideal) in P;
2. $P \setminus I$ is a poset coideal (ideal) of P;
3. $\widetilde{I}$ is a poset coideal (ideal) of $\widetilde{P}$.

Let P be an arbitrary finite poset and write $\mathscr{J}(P)$ for the poset which consists of all poset ideals of P ordered by inclusion.

For example, if P is an antichain, i.e., any two elements of P are incomparable, then $\mathscr{J}(P) \cong \mathscr{B}_P$, where $\mathscr{B}_P$ is the Boolean lattice consisting of all subsets of P. The rank of $\mathscr{B}_P$ is the cardinality of P.

Since the union $I \cup J$ and the intersection $I \cap J$ of poset ideals I and J of P are also poset ideals of P, the poset $\mathscr{J}(P)$ is in fact a lattice.

We say that a poset P is *isomorphic* to a poset Q if there exists a bijection $\theta : P \to Q$ such that $\alpha \le \beta$ in P if and only if $\theta(\alpha) \le \theta(\beta)$ in Q.

The most important class of lattices from the combinatorial point of view is the distributive lattice. And one of the most influential results in the classical lattice theory is Birkhoff's fundamental structure theorem for the finite distributive lattice.

Theorem 1.2 (Birkhoff)**.** *Let $\mathscr{L}$ be a finite distributive lattice. Then there exists a unique (up to isomorphism) poset P such that $\mathscr{L}$ is isomorphic to $\mathscr{J}(P)$.*

One finds the proof, for example, in [9, Theorem 3.4.1]. In fact, P can be chosen as the join irreducible subposet of $\mathscr{L}$.

The outline of the proof is as follows: Let $\mathscr{L}$ be a finite lattice, and let P be the set of join irreducible elements of $\mathscr{L}$. As in [8] we associate to each element $p \in \mathscr{L}$ the poset ideal $\ell(p) = \{q \in P \colon q \le p\}$ of P. This establishes a map $\ell \colon \mathscr{L} \to \mathscr{J}(P)$, which we call the *canonical embedding* into the distributive lattice $\mathscr{J}(P)$. Note that ℓ is an isomorphism if and only if $\mathscr{L}$ is distributive.

We call the cardinality of $\ell(p)$ the *degree* of p, and denote it by $\deg p$.

The map ℓ has the following properties.

Lemma 1.3. *Let $\mathscr{L}$ be a finite lattice, ℓ the canonical embedding and $s, t \in \mathscr{L}$ any two elements. We have*

1. *$s = t$ if and only if $\ell(s) = \ell(t)$;*
2. *$s \le t$ if and only if $\ell(s) \subseteq \ell(t)$;*
3. *$\ell(s) \cap \ell(t) = \ell(s \wedge t)$.*

This lemma implies in particular that ℓ is an injective order preserving map. In general however, ℓ is not an embedding of lattices.

As a consequence of Remark 1.1 we have the following

Lemma 1.4. $\widetilde{\mathscr{L}} \cong \mathscr{J}(\widetilde{P})$.

Proof. Let $q \in \widetilde{\mathscr{L}}$. Since the underlying set of $\widetilde{\mathscr{L}}$ is the same as that of $\mathscr{L}$, we may apply $\ell \colon \mathscr{L} \to \mathscr{J}(P)$ to q. Then $\widetilde{\mathscr{L}} \to \mathscr{J}(\widetilde{P})$, $q \mapsto P \setminus \ell(q)$ is the desired isomorphism. $\square$

We now introduce the squarefree monomial ideal $H_{\mathscr{L}}$ associated with a finite lattice $\mathscr{L}$. Let K be a field and $S = K[\{x_p, y_p\}_{p \in P}]$ the polynomial ring in $2|P|$ variables over K.

For each element $q \in \mathscr{L}$ write

$$u_q = \prod_{p \in \ell(q)} x_p \prod_{p \in P \setminus \ell(q)} y_p,$$

and set $H_{\mathscr{L}} = (u_q)_{q \in \mathscr{L}}$. We call $H_{\mathscr{L}}$ the *Hibi ideal* of $\mathscr{L}$. It is easy to see that the height of $H_{\mathscr{L}}$ is 2.

Recall that a finite lattice $\mathscr{L}$ is *upper semimodular* if $\mathscr{L}$ satisfies either of the following two conditions.

(1) $\mathscr{L}$ is graded, and the rank function ρ of $\mathscr{L}$ satisfies $\rho(\alpha) + \rho(\beta) \geq \rho(\alpha \wedge \beta) + \rho(\alpha \vee \beta)$ for all $\alpha, \beta \in \mathscr{L}$.
(2) If α and β both cover $\alpha \wedge \beta$, then $\alpha \vee \beta$ covers both α and β.

Among the upper semimodular lattices the distributive lattices can be characterized algebraically. For this we need following concept: let I be a monomial ideal in a polynomial ring with the (unique) minimal set $G(I)$ of monomial generators. The ideal I is called a *linear quotient ideal* if the elements of $G(I)$ can be ordered $u_1, \ldots, u_m$ such that the colon ideals $(u_1, \ldots, u_{i-1}) : u_i$ are generated by variables. If I is squarefree, then $I = (u_1, \ldots, u_m)$ is a linear quotient ideal (in this order) if and only if for each i and each $j < i$ there exists $k < i$ such that $u_k/[u_k, u_i]$ is a variable which divides u_j. Here $[u, v]$ denotes the greatest common divisor of u and v.

It is easy to see that a linear quotient ideal I has a linear resolution, if all generators of I have the same degree. If I is an ideal with linear resolution, then all generators of I have the same degree, say d. In this case we also say I has a d-linear resolution.

The following characterization of finite distributive lattices is an immediate consequence of [8, Theorem 1.3].

Proposition 1.5. *Let $\mathscr{L}$ be an arbitrary finite upper semimodular lattice. The following conditions are equivalent:*

(1) *$\mathscr{L}$ is distributive;*
(2) *$H_{\mathscr{L}}$ has linear quotients;*
(3) *$H_{\mathscr{L}}$ has a linear resolution;*
(4) *$H_{\mathscr{L}}$ has linear relations.*

Now we recall some concepts related to simplicial complex and fix some notation. Let Δ be a simplicial complex on the vertex set $[n] = \{1, \ldots, n\}$, $R = K[x_1, \ldots, x_n]$ the polynomial ring in n variables over a field K. We denote $\mathscr{F}(\Delta)$ the set of facets (maximal faces) of Δ. The simplicial complex

$$\Delta^{\vee} = \{[n] \setminus F : F \notin \Delta\}$$

is called the *Alexander dual* of Δ. One has $(\Delta^{\vee})^{\vee} = \Delta$.

A *vertex cover* of Δ is a set $G \subset [n]$ such that $G \cap F \neq \emptyset$ for all $F \in \mathscr{F}(\Delta)$. We say a vertex cover G of Δ is *minimal*, if each proper subset of G is not a vertex cover of Δ. We denote by $\mathscr{C}(\Delta)$ the set of minimal vertex covers of Δ. If all the minimal vertex cover of Δ have the same cardinality, then we say Δ is *unmixed*. For $F = \{i_1, \ldots, i_k\} \subset [n]$, let P_F be the prime ideal generated by $x_{i_1}, \ldots, x_{i_k}$, and set $F^c = [n] \setminus F$. As usual we denote by I_{Δ} the Stanley–Reisner ideal and $K[\Delta] = R/I_{\Delta}$ the Stanley–Reisner ring of Δ. The following proposition is an elementary but important property of the Stanley–Reisner ideal. One finds the proof for example in [2, Theorem 5.1.4].

Proposition 1.6. *Let Δ be a simplicial complex over the vertex set $[n]$. Then*

$$I_\Delta = \bigcap_{F \in \mathscr{F}(\Delta)} P_{F^c}.$$

The *facet ideal* is defined to be

$$I(\Delta) = (x_F : F \in \mathscr{F}(\Delta)),$$

where $x_F = \prod_{i \in F} x_i$.

Let Γ be the unique simplicial complex such that $I_\Delta = I(\Gamma)$. Then

$$I_\Delta = \bigcap_{F \in \mathscr{C}(\Gamma)} P_F \quad \text{and} \quad I_{\Delta^\vee} = (x_F : F \in \mathscr{C}(\Gamma)). \tag{1}$$

Set $\Delta^c = \langle F^c : F \in \mathscr{F}(\Delta) \rangle$. Then

$$I_{\Delta^\vee} = I(\Delta^c). \tag{2}$$

The easy proofs can be found for example in [7].

The proof of the following simple lemma can be found for example in [4, Proposition 1.8].

Lemma 1.7. *Let Δ be a simplicial complex on the vertex set $[n]$ and $I(\Delta)$ the facet ideal of Δ. Then an ideal $P = (x_{i_1}, \ldots, x_{i_s})$ is a minimal prime of $I(\Delta)$ if and only if $\{i_1, \ldots, i_s\}$ is a minimal vertex cover of Δ.*

Let Δ be a simplicial complex and Γ the unique simplicial complex with $I_\Gamma = I(\Delta)$. By using Proposition 1.6 and the previous lemma, we have:

Corollary 1.8. *A subset F of $[n]$ is a facet of Γ if and only if F^c is a minimal vertex cover of Δ.*

We say an ideal I in a ring R is *Cohen–Macaulay* if R/I is a Cohen–Macaulay R-module. Let Δ be a simplicial complex such that $I(\Delta)$ is a Cohen–Macaulay ideal. Since any Cohen–Macaulay simplicial complex is pure, using Corollary1.8, we have Δ is unmixed.

Let I be a squarefree monomial ideal. Then $I = I_\Delta$ for some simplicial complex Δ. For the convenience we write I^* for $I_{\Delta^\vee}$.

Lemma 1.9. *Let I and J be two squarefree monomial ideals. Then*

$$(I \cap J)^* = I^* + J^*.$$

Proof. Let P be a monomial prime ideal in R. Then $I \cap J \subseteq P$ if and only if $I \subseteq P$ or $J \subseteq P$. The assertion follows from (1). ☐

The following theorem gives important algebraic properties of Alexander duality.

Theorem 1.10. *Let K be a field, Δ a simplicial complex, I_Δ the Stanley–Reisner ideal and $K[\Delta]$ the Stanley–Reisner ring of Δ. Then*

1. (Eagon–Reiner [3]) $K[\Delta]$ *is Cohen–Macaulay* $\Longleftrightarrow$ $I_{\Delta^\vee}$ *has a linear resolution.*
2. (Herzog–Hibi–Zheng [7]) Δ *is shellable* $\Longleftrightarrow$ $I_{\Delta^\vee}$ *has linear quotients.*

2. A CLASS OF UNMIXED SIMPLICIAL COMPLEXES

A simplicial complex Δ on the vertex set $[n]$ is *Cohen–Macaulay* over a field K, if the Stanley–Reisner ideal I_Δ of Δ is a Cohen–Macaulay ideal, while for a graph G, we say G is Cohen–Macaulay, if the edge ideal $I(G)$ of G is a Cohen–Macaulay ideal.

A graph G is *bipartite* if its vertex set V can be partitioned into disjoint subsets V_1 and V_2 such that every edge $\{v_1,v_2\}$ of G satisfies $v_1 \in V_1$ and $v_2 \in V_2$. Let G be a bipartite graph with no isolated vertex on the vertex set $V \cup V'$, where $V \cap V' = \emptyset$ and $|V| = |V'|$. In [6, Theorem 2.4], the authors showed that a bipartite graph G is a Cohen–Macaulay if and only if $I(G) = H^*_{\mathscr{L}}$ for some distributive lattice $\mathscr{L}$. Later in [8], the authors considered simplicial complexes Δ on the vertex set $V \cup V'$ with $V \cap V' = \emptyset$ and $|V| = |V'|$, such that

(1) there is no $F \in \mathscr{F}(\Delta)$ with $F \subset V$, and
(2) $G = \{F \in \mathscr{F}(\Delta)\colon F \cap V \neq \emptyset, \quad F \cap V' \neq \emptyset\}$ is a Cohen–Macaulay bipartite graph with no isolated vertex,

and showed when the facet ideal $I(\Delta)$ of Δ is Cohen–Macaulay, see [8, Theorem 4.3].

In this section we will consider a further generalization of Theorem 2.4 in [6]. For this we need some preparation.

The poset ideals and poset coideals of lattices are special subsets of lattices. Now we introduce a more general class of subsets of lattices:

Definition 2.1. Let $\mathscr{L}$ be a lattice. A subset $\mathscr{S}$ of $\mathscr{L}$ is called a *segment* of $\mathscr{L}$, if for all $p,q \in \mathscr{S}$ with $p \leq q$, we have $[p,q] \subseteq \mathscr{S}$.

It is clear that any poset ideal and any poset coideal of a lattice $\mathscr{L}$ are segments of $\mathscr{L}$. Furthermore, we have

Lemma 2.2. *Let $\mathscr{L}$ be a lattice, $\mathscr{S}$ a subset of $\mathscr{L}$. Then the following statements are equivalent:*

(1) *$\mathscr{S}$ is a segment of $\mathscr{L}$;*
(2) *$\mathscr{S}$ is the intersection of a poset ideal and a poset coideal of $\mathscr{L}$.*

Proof. (1) $\Rightarrow$ (2): Let $\mathscr{I} = \{r \in \mathscr{L} :$ there exists an element $s \in \mathscr{S}$ such that $r \leq s\}$ and $\mathscr{J} = \{r \in \mathscr{L} :$ there exists an element $s \in \mathscr{S}$ such that $r \geq s\}$. Then $\mathscr{I}$ is a poset ideal of $\mathscr{L}$ and $\mathscr{J}$ is a poset coideal of $\mathscr{L}$. For any $s \in \mathscr{S}$, we have $s \in \mathscr{I} \cap \mathscr{J}$. This implies $\mathscr{S} \subseteq \mathscr{I} \cap \mathscr{J}$. Now let r be an arbitrary element in $\mathscr{I} \cap \mathscr{J}$. Then there exist $p,q \in \mathscr{S}$ such that $p \leq r \leq q$, i.e., $r \in [p,q]$. Since $\mathscr{S}$ is a segment, we have $r \in \mathscr{S}$. Hence $\mathscr{I} \cap \mathscr{J} \subseteq \mathscr{S}$.

(2) $\Rightarrow$ (1): Assume $\mathscr{S} = \mathscr{I} \cap \mathscr{J}$, where $\mathscr{I}$ is a poset ideal of $\mathscr{L}$ and $\mathscr{J}$ is a poset coideal of $\mathscr{L}$. Let $r \in [p,q]$ with $p,q \in \mathscr{S}$ and $p \leq q$. Since $q \in \mathscr{I}$ and $r \leq q$, we have $r \in \mathscr{I}$. Since $p \in \mathscr{J}$ and $r \geq p$, we have $r \in \mathscr{J}$. Hence $r \in \mathscr{I} \cap \mathscr{J} = \mathscr{S}$. This implies that $\mathscr{S}$ is a segment of $\mathscr{L}$. ☐

Remark 2.3. Let $\mathscr{S}$ be a segment of a lattice $\mathscr{L}$. The poset ideal $\mathscr{I}$ and poset coideal $\mathscr{J}$ with the property $\mathscr{S} = \mathscr{I} \cap \mathscr{J}$ are not uniquely determined. The poset ideal $\mathscr{I}$ and poset coideal $\mathscr{J}$ in the proof (1) $\Rightarrow$ (2) of Lemma 2.2 are the minimal one with this property.

Let G be a Cohen–Macaulay bipartite graph on the vertex set $V \cup V'$ with $V \cap V' = \emptyset$ and $|V| = |V'| = n$, and $S = K[x_1,\ldots,x_n,y_1,\ldots,y_n]$ the polynomial ring over a field K.

Recall from [6, Theorem 2.4] that the vertices $V = \{x_1, \dots, x_n\}$ and $V' = \{y_1, \dots, y_n\}$ can be labeled such that there exists a partial order $<$ on V with the property that $\{x_i, y_j\}$ is an edge of G if and only if $x_i \le x_j$. Moreover it is shown that for $P = (V, <)$ the distributive lattice $\mathscr{J}(P)$ satisfies $H^*_{\mathscr{J}(P)} = I(G)$. We denote this lattice by $\mathscr{L}(G)$. As a generalization of this result we have:

Theorem 2.4. *Let Δ be a simplicial complex on the vertex set $V \cup V'$ with $V \cap V' = \emptyset$ and $|V| = |V'|$. Suppose that $G = \{F \in \mathscr{F}(\Delta) : F \cap V \neq \emptyset, \quad F \cap V' \neq \emptyset\}$ is a Cohen–Macaulay bipartite graph with no isolated vertex. Then the following conditions are equivalent:*

(1) *Δ is unmixed, and all minimal vertex covers of Δ have cardinality $|V|$;*
(2) *there exists a lattice segment $\mathscr{S} \subseteq \mathscr{L}(G)$ such that $H^*_{\mathscr{S}} = I(\Delta)$.*

Proof. (1) $\Rightarrow$ (2): Let Γ be the (unique) simplicial complex defined by the equation $I_\Gamma = I(\Delta)$. Since Δ is unmixed, we have Γ is pure. Let $V = \{x_1, \dots, x_n\}$ and $V' = \{y_1, \dots, y_n\}$ with the labeling as described before this theorem. Since Δ is a complex with $2n$ vertices and the minimal vertex cover of Δ has cardinality n, it follows from Corollary 1.8, that $|F| = n$ for each $F \in \mathscr{F}(\Gamma)$.

Let Γ_0 be the simplicial complex on $V \cup V'$ with $I_{\Gamma_0} = I(G)$. Then any minimal vertex cover of Δ is a minimal vertex cover of G. Indeed, a minimal vertex cover C of Δ is also a vertex cover of G, and it has cardinality n, by assumption. On the other hand, since G contains all the edges $\{x_i, y_i\}$, each vertex cover of G has at least cardinality n. Hence C is a minimal vertex cover of G.

It follows that each facet of Γ is a facet of Γ_0. In other words, each minimal nonface of $\Gamma^\vee$ is a minimal nonface of $\Gamma_0^\vee$. Therefore, $G(I_{\Gamma^\vee}) \subset G(I_{\Gamma_0^\vee}) = G(H_{\mathscr{L}(G)})$. That is, there exists a subset $\mathscr{S} \neq \emptyset$ of $\mathscr{L}(G)$, such that $G(I_{\Gamma^\vee}) = \{u_s : s \in \mathscr{S}\}$, and this implies that $I(\Delta) = H^*_{\mathscr{S}}$.

Now, what we must prove is that for any $p, q \in \mathscr{S}$ with $p \le q$ one has $[p, q] \subseteq \mathscr{S}$. Suppose, on the contrary, there exist two elements δ and ξ of $\mathscr{L}(G)$ with $\xi < \delta$, and $\gamma \in \mathscr{L}(G)$ such that $\gamma \in [\xi, \delta]$ but $\gamma \notin \mathscr{S}$.

Recall that the elements of $\mathscr{L}(G)$ are poset ideals of $P = (V, <)$. To simplify the notation, we will assume that $\xi = \{x_1, \dots, x_l\}$, $\gamma = \{x_1, \dots, x_r\}$ and $\delta = \{x_1, \dots, x_k\}$ with $l < r < k$. Since $\xi = \{x_1, \dots, x_l\} \in \mathscr{S}$, we have $x_1 \cdots x_l y_{l+1} \cdots y_n \in G(H_{\mathscr{S}})$. Thus $\{x_1, \dots, x_l, y_{l+1}, \dots, y_n\}$ is a minimal vertex cover of Δ. It follows from Corollary 1.8 that $\{y_1, \dots, y_l, x_{l+1}, \dots, x_n\} \in \mathscr{F}(\Gamma)$. By the same reason we have $\{y_1, \dots, y_k, x_{k+1}, \dots, x_n\} \in \mathscr{F}(\Gamma)$, but $\{y_1, \dots, y_r, x_{r+1}, \dots, x_n\} \notin \mathscr{F}(\Gamma)$. Hence there exists a monomial generator u of $I_\Gamma = I(\Delta)$ such that u does not divide $y_1 \cdots y_k x_{k+1} \cdots x_n$ and $y_1 \cdots y_l x_{l+1} \cdots x_n$, but divides $y_1 \cdots y_r x_{r+1} \cdots x_n$. Hence there exists an i with $r < i \le k$, such that $x_i \mid u$ and a j with $l < j \le r$ such that $y_j \mid u$. By our assumption, $u = x_i y_j$. By our labeling of the vertices it follows that $x_i < x_j$ in P. Since $j \le r$, we have that $x_j \in \gamma$. Since γ is a poset ideal it follows that also $x_i \in \gamma$. This is impossible, since $i > r$.

(2) $\Rightarrow$ (1): Since all generators of $H_{\mathscr{S}}$ are of same degree and since $H^*_{\mathscr{S}} = I(\Delta)$, it follows that Δ is unmixed. $\square$

The following two simplicial complexes are unmixed and satisfy the assumption in Theorem 2.4 with $V_\Delta = \{a, b, c, d\}$, $V'_\Delta = \{u, v, w, x\}$, and $V_{\Delta'} = \{a, b, c\}$, $V'_{\Delta'} = \{u, v, w\}$.





The segments $\mathscr{S}$ and $\mathscr{S}'$ such that $I(\Delta) = H^*_{\mathscr{S}}$ and $I(\Delta') = H^*_{\mathscr{S}'}$ are given in the next figures. For the simplicity, sets in these figures are written as monomials. For example $abcd$ stands for $\{a,b,c,d\}$. The elements of the segments are indicated by the bullet vertices.





Note that the facet ideal $I(\Delta)$ of δ is Cohen–Macaulay, while the facet ideal $I(\Delta')$ of Δ' is not. This is because the ideal $H_{\mathscr{S}} = \{avwx, buwx, cuvx, abwx, acvx, bcux, abcx, bcdu\}$ has a linear resolution, while the ideal $H_{\mathscr{S}'} = \{avw, buw, cuv, abw, acv, bcu\}$ has no linear resolution. It is therefore of interest to know for which kind of segments $\mathscr{S}$ of a finite distributive lattice $\mathscr{L}$, the ideal $H_{\mathscr{S}}$ has a linear resolution.

3. LATTICE SEGMENTS AND POSET IDEALS

We use the notation as in the previous sections. We have already seen that $\mathscr{S} = \mathscr{I} \cap \mathscr{J}$ where $\mathscr{I}$ is a poset ideal and $\mathscr{J}$ a poset coideal in $\mathscr{L}$. In case $H_{\mathscr{S}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$, necessary and sufficient conditions for $H_{\mathscr{S}}$ to have a linear resolution will be given. We will also discuss when $H_{\mathscr{S}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$.

Let $p \in \mathscr{L}$, and set $N(p)$ for the set of lower neighbors, and $M(p)$ for the set of upper neighbors of p.

Let P be the set of join-irreducible elements of $\mathscr{L}$, and $<$ a total order on $\mathscr{L}$ which extends the partial order on P. For a subset $T \subset P$ and $q \in P$ we set

$$\lambda(q;T) = |\{r \in T : r < q\}|.$$

For each element $q \in N(p)$, we have $|\ell(p) \setminus \ell(q)| = 1$. We denote the unique element in $\ell(p) \setminus \ell(q)$ by $p \setminus q$. Let $p \in \mathscr{L}$, $S \subseteq N(p)$ and $T \subseteq M(p)$. We set $p \setminus S = \{p \setminus s : s \in S\}$, and $T \setminus p = \{t \setminus p : t \in T\}$. Note that $p \setminus S$ and $T \setminus p$ both are subset of P. Let $p \in \mathscr{L}$ and $S \subset \mathscr{L}$. We also set $p \vee S = \{p \vee s : s \in S\}$ and $p \wedge S = \{p \wedge s : s \in S\}$. The following theorem is shown in [8]:

Theorem 3.1 (Herzog-Hibi-Zheng [8])**.** *Let $\mathscr{L}$ be finite meet-semilattice.*

1. *There exists a finite multigraded free S-resolution $\mathbb{F}$ of $H_{\mathscr{L}}$ such that for each $i \geq 0$, the free module $\mathbb{F}_i$ has a basis with basis elements*
$$b(p;S)$$
where $p \in \mathscr{L}$ and $S \subseteq N(p)$ with $|S| = i$. The multidegree of $b(p;S)$ is the least common multiple of u_p and all monomials u_q with $q \in S$.
2. *The following conditions are equivalent:*
   - (a) *the resolution constructed in* (1) *is minimal;*
   - (b) *for any $p \in \mathscr{L}$ and for any proper subset $S \subset N(p)$ the meet $\bigwedge\{q\colon q \in S\}$ is strictly greater than the meet $\bigwedge\{q\colon q \in N(p)\}$.*
3. *If $\mathbb{F}$ is minimal, then the differential ∂ in $\mathbb{F}$ is as follows: for each $p \in \mathscr{L}$ and each $S \subset N(p)$, one has*
$$\partial(b(p;S)) = \sum_{q \in S} (-1)^{\lambda(p\setminus q; p\setminus S)} (y_{p\setminus q} b(p; S \setminus \{q\}) - x_{p\setminus q} b(q; q \wedge (S \setminus \{q\})).$$

Corollary 3.2. *Let $\mathscr{L}$ be a finite distributive lattice and $\mathscr{I}$ a poset ideal (coideal) of $\mathscr{L}$. Then the minimal free resolution of $H_{\mathscr{I}}$ is linear.*

Proof. Let $\mathscr{I}$ be a poset ideal of $\mathscr{L}$. Then $\mathscr{I}$ is a meet-semilattice, and has property (2)(b) of Theorem 3.1. Hence the free resolution of $H_{\mathscr{I}}$ as described in Theorem 3.1(1) is minimal. For any $p \in \mathscr{I}$ and any $S \subseteq N(p)$, the total degree of $b(p;S)$ equals $\operatorname{rank} \mathscr{L} + |S|$. This shows that the resolution of $H_{\mathscr{I}}$ is linear.

Now assume that $\mathscr{I}$ is a poset coideal. Then by Remark 1.1, $\widetilde{\mathscr{I}}$ is a poset ideal in $\widetilde{\mathscr{L}}$. Therefore $H_{\widetilde{\mathscr{I}}}$ has a linear resolution by the first part of the proof. By Lemma 1.4 (and its proof) the canonical labeling $\widetilde{\ell}$ of $\widetilde{\mathscr{L}}$ is given by $\widetilde{\ell}(p) = P \setminus \ell(p)$ for all $p \in \widetilde{\mathscr{L}}$. It follows that $H_{\widetilde{\mathscr{I}}}$ is generated by the monomials $\widetilde{u}_p = x_{\widetilde{\ell}(p)} y_{P\setminus\widetilde{\ell}(p)} = x_{P\setminus\ell(p)} y_{\ell(p)}$. Now we apply the following involution

$$\sigma\colon K[\{x_p, y_p\}_{p\in P}] \to K[\{x_p, y_p\}_{p\in P}], \quad x_p \mapsto y_p \quad \text{and} \quad y_p \mapsto x_p, \tag{3}$$

and we obtain $\sigma(H_{\widetilde{\mathscr{I}}}) = H_{\mathscr{I}}$. This shows that $H_{\mathscr{I}}$ has a linear resolution, too. $\square$

As we have already seen that for a poset ideal $\mathscr{I}$ and a poset coideal $\mathscr{J}$ of a finite distributive lattice $\mathscr{L}$, the ideal $H_{\mathscr{I}}$ and $H_{\mathscr{J}}$ both have linear resolutions, one might except that if we write $\mathscr{S} = \mathscr{I} \cap \mathscr{J}$ for some poset ideal $\mathscr{I}$ and some poset coideal $\mathscr{J}$ of $\mathscr{L}$ and if $H_{\mathscr{S}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$, then the ideal $H_{\mathscr{S}}$ has a linear resolution. However there are two questions arising: (1) when $H_{\mathscr{I}\cap\mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$ and (2) whether $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has a linear resolution. In general, the intersection of two ideals with linear resolutions need not to have a linear resolution, even for the special ideals $H_{\mathscr{I}}$ and $H_{\mathscr{J}}$. For example, consider the Boolean lattice $\mathscr{B}_3$ of rank 3, and let $\mathscr{I} = \mathscr{B}_3 \setminus \{\hat{1}\}$ and $\mathscr{J} = \mathscr{B}_3 \setminus \{\hat{0}\}$. Then $H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$, but it has no linear resolution.

To see when $H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$ and when $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has a linear resolution, we need some preparation.

Let $\mathscr{I}$ be a poset ideal of a finite distributive lattice $\mathscr{L}$ and let $\mathbb{F}$ be the minimal free resolution of $H_{\mathscr{L}}$ and $\mathbb{P}$ the minimal free resolution of $H_{\mathscr{I}}$. Then by Theorem 3.1 one sees that $\mathbb{P}$ is a subcomplex of $\mathbb{F}$. More precisely we have

Lemma 3.3. *For each $i \geq 0$ there exists an injective map $\operatorname{Tor}_i(K, H_{\mathscr{I}}) \to \operatorname{Tor}_i(K, H_{\mathscr{L}})$ which maps the basis elements $b(p;S)$ of $\operatorname{Tor}_i(K, H_{\mathscr{I}})$ to the corresponding basis elements of $\operatorname{Tor}_i(K, H_{\mathscr{L}})$.*

For convenience, in this lemma and the remaining of this section the basis elements in a free resolution and corresponding basis in the Tor-groups are denoted by the same symbol.

Let $\mathscr{J}$ be a poset coideal of $\mathscr{L}$. Then $\widetilde{\mathscr{J}}$ is a poset ideal of $\widetilde{\mathscr{L}}$. Let $\widetilde{\mathbb{F}}$ be the minimal multigraded free resolution of $H_{\widetilde{\mathscr{L}}}$ and $\widetilde{\mathbb{T}}$ the minimal multigraded free resolution of $H_{\widetilde{\mathscr{J}}}$, as described in Theorem 3.1. Then $\widetilde{\mathbb{T}}$ is a subcomplex of $\widetilde{\mathbb{F}}$, and the injective map $\operatorname{Tor}_i(K, H_{\widetilde{\mathscr{J}}}) \to \operatorname{Tor}_i(K, H_{\widetilde{\mathscr{L}}})$ is as described in Lemma 3.3. Since $\sigma(H_{\widetilde{\mathscr{L}}}) = H_{\mathscr{L}}$, we have $\sigma(\widetilde{\mathbb{F}})$ is a minimal multigraded free resolution of $H_{\mathscr{L}}$. Since $\mathbb{F}$ is also a minimal multigraded free resolution of $H_{\mathscr{L}}$, it is natural to ask what is the isomorphic chain map from $\widetilde{\mathbb{F}}$ to $\mathbb{F}$.

To answer this question we need the following two lemmata:

Lemma 3.4. *Let $\widetilde{\mathscr{L}}$ be the dual of the distributive lattice $\mathscr{L}$ and $\widetilde{\mathbb{F}}$ the minimal multigraded free resolution of $H_{\widetilde{\mathscr{L}}}$ as in Theorem 3.1. Then*

(1) *for each $i \geq 0$, the free module $\widetilde{F}_i$ has a basis with basis elements $\widetilde{b}(r;T)$ with $r \in \mathscr{L}$, $T \subseteq M(r)$ in $\mathscr{L}$ and $|T| = i$. The multidegree of $\widetilde{b}(r;T)$ is the least common multiple of $\widetilde{u}_r$ and all monomials $\widetilde{u}_s$ with $s \in T$;*

(2) *the differential $\widetilde{\partial}$ in $\widetilde{\mathbb{F}}$ is as follows: for each $r \in \mathscr{L}$ and each $T \subseteq M(r)$, one has*

$$\widetilde{\partial}(\widetilde{b}(r;T)) = \sum_{s \in T} (-1)^{\lambda(s \setminus r; T \setminus r)} (y_{s \setminus r} \widetilde{b}(r; T \setminus \{s\}) - x_{s \setminus r} \widetilde{b}(s; s \vee (T \setminus \{s\}))).$$

Proof. We may assume that $\widetilde{\mathbb{F}}$ is a minimal free resolution of $H_{\widetilde{\mathscr{L}}}$ as described in Theorem 3.1. Therefore $\widetilde{\mathbb{F}}$ has a basis $\widetilde{b}(r;T)$ where $r \in \widetilde{\mathscr{L}}$ and T is a subset of lower neighbors of r in $\widetilde{\mathscr{L}}$. Moreover, we have

$$\widetilde{\partial}(\widetilde{b}(r;T)) = \sum_{s \in T} (-1)^{\lambda(\widetilde{r \setminus s}; \widetilde{r \setminus T})} (y_{\widetilde{\ell}(r) \setminus \widetilde{\ell}(s)} \widetilde{b}(r; T \setminus \{s\}) - x_{\widetilde{\ell}(r) \setminus \widetilde{\ell}(s)} \widetilde{b}(s; s \wedge (T \setminus \{s\}))),$$

where $\widetilde{r \setminus s}$ denote the unique element in $\widetilde{\ell}(r) \setminus \widetilde{\ell}(s)$ and $\widetilde{r \setminus T}$ the set $\{\widetilde{\ell}(r) \setminus \widetilde{\ell}(s) : s \in T\}$.

Notice that for any element $r \in \widetilde{\mathscr{L}}$ (hence $r \in \mathscr{L}$, too), a lower (upper) neighbor of r in $\widetilde{\mathscr{L}}$ is just a upper (lower) neighbor of r in $\mathscr{L}$, and for any two element p and q in $\widetilde{\mathscr{L}}$, the meet (join) of p and q in $\widetilde{\mathscr{L}}$ is just the join (meet) of them in $\mathscr{L}$.

Let $r \in \widetilde{\mathscr{L}}$ and s a lower neighbor of r in $\widetilde{\mathscr{L}}$. We have

$$\widetilde{\ell}(r) \setminus \widetilde{\ell}(s) = (P \setminus \ell(r)) \setminus (P \setminus \ell(s)) = s \setminus r$$

and

$$\lambda(\widetilde{r \setminus s}; \widetilde{r \setminus T}) = \lambda(\widetilde{\ell}(r) \setminus \widetilde{\ell}(s); \{\widetilde{\ell}(r) \setminus \widetilde{\ell}(s) : s \in T\}) = \lambda(s \setminus r; \{s \setminus r : s \in T\}) = \lambda(s \setminus r; T \setminus r).$$

Thus we obtain the desired formula. □

Let S be any subset of $\mathscr{L}$. We set $\vee S = \vee\{s : s \in S\}$ and $\wedge S = \wedge\{s : s \in S\}$.

Lemma 3.5. *Let $\mathscr{L}$ be a finite distributive lattice, $p \in \mathscr{L}$ and $S \subseteq N(p)$ with $|S| = i$. Let $r = \wedge\{q : q \in S\}$, and T the set of all upper neighbors of r in the interval $[r, p]$. Then*

1. *$|T| = i$ and $\vee T = p$;*
2. *$\mathrm{lcm}(u_p, \{u_q\}_{q \in S}) = \mathrm{lcm}(u_r, \{u_s\}_{s \in T})$;*
3. *for any $r' \neq r$ and $T' \subseteq M(r')$, one has $\mathrm{lcm}(u_{r'}, \{u_{s'}\}_{s' \in T'}) \neq \mathrm{lcm}(u_p, \{u_q\}_{q \in S})$.*

Proof. (1) Since $\mathscr{L}$ is a distributive lattice, the interval $[r, p]$ is a Boolean lattice. Hence $|T| = |S| = i$ and $\vee T = p$.

(2) The monomial associated to p is $u_p = x_{\ell(p)} y_{P \setminus \ell(p)}$, where P is the set of join irreducible elements of $\mathscr{L}$. Let $q \in N(p)$. Then $u_q = x_{\ell(p) \setminus (p \setminus q)} y_{(P \setminus \ell(p)) \cup (p \setminus q)}$. Hence

$$\mathrm{lcm}(u_p, \{u_q\}_{q \in S}) = x_{\ell(p)} y_{(P \setminus \ell(p)) \cup (\bigcup_{q \in S} (p \setminus q))}.$$

On the other hand, $\ell(r) = \ell(p) \setminus (\bigcup_{q \in S} (p \setminus q))$, $u_r = x_{\ell(p) \setminus (\bigcup_{q \in S} (p \setminus q))} y_{P \setminus (\ell(p) \setminus \bigcup_{q \in S} (p \setminus q))}$. Since $\ell(p) = \ell(r) \cup (\bigcup_{s \in T} \ell(s))$, we have

$$\mathrm{lcm}(u_r, \{u_s\}_{s \in T}) = x_{\ell(p)} y_{P \setminus (\ell(p) \setminus (\bigcup_{q \in S} (p \setminus q)))}.$$

Since $\ell(p) \subseteq P$ and $\bigcup_{p \in S} (p \setminus q) \subseteq P$, we have

$$(P \setminus \ell(p)) \cup (\bigcup_{q \in S} (p \setminus q)) = P \setminus (\ell(p) \setminus \bigcup_{q \in S} (p \setminus q)).$$

Hence (2) follows.

(3) As in the proof of (2) we see that the y-part of $\mathrm{lcm}(u_{r'}, \{u_{s'}\}_{s' \in T'})$ equals the y-part of $u_{r'}$. Since for any $r' \neq r$, we have $\ell(r') \neq \ell(r)$. The assertion follows from (2). □

We fix some notation. For each element $r \in \mathscr{L}$ and $T \subseteq M(r)$, we write r^T for the join of all elements in T, and T_r the set of all lower neighbors of r^T in the interval $[r, r^T]$.

The polynomial ring S viewed as a S-module via the involution $\sigma : S \to S$ is denoted by ${}^{\sigma}S$. Let $\widetilde{\mathbb{F}}$ be the minimal free resolution of the ideal $H_{\widetilde{\mathscr{L}}}$ with basis elements $\widetilde{b}(r; T)$ as described in Lemma 3.4. Then $\widetilde{\mathbb{F}} \bigotimes_S {}^{\sigma}S$ with basis elements $\widetilde{b}(r; T) \bigotimes_S 1$ is a minimal free resolution of $\sigma(H_{\widetilde{\mathscr{L}}}) = H_{\mathscr{L}}$. We denote the complex $\widetilde{\mathbb{F}} \bigotimes_S {}^{\sigma}S$ by $\sigma(\widetilde{\mathbb{F}})$ and the basis elements $\widetilde{b}(r; T) \bigotimes_S 1$ by $\sigma(\widetilde{b}(r; T))$.

Proposition 3.6. *Let $\mathscr{L}$ be a finite distributive lattice, and let $\mathbb{F}$ and $\widetilde{\mathbb{F}}$ be the minimal multigraded free resolutions of $H_{\mathscr{L}}$ and $H_{\widetilde{\mathscr{L}}}$, respectively. Then the map $\pi : \sigma(\widetilde{\mathbb{F}}) \to \mathbb{F}$ with $\pi(\sigma(\widetilde{b}(r; T))) = (-1)^{|T|} b(r^T; T_r)$ is an isomorphism of complexes.*

Proof. Let $s \in T$ and $q = \vee\{s' \in T : s' \neq s\}$. Let $|T| = i$. By Lemma 3.4, we have

$$
\begin{aligned}
&(4)\pi_{i-1}(\widetilde{\partial}_i(\sigma(\widetilde{b}(r;T))))\\
= \quad & \pi_{i-1}(\sum_{s\in T}(-1)^{\lambda(s\setminus r;T\setminus r)}(y_{s\setminus r}\widetilde{b}(r;T\setminus\{s\}) - x_{s\setminus r}\widetilde{b}(s;s\vee(T\setminus\{s\}))))\\
= \quad & \sum_{s\in T}(-1)^{\lambda(s\setminus r;T\setminus r)+(i-1)+1}(y_{s\setminus r}b(s^{s\vee(T\setminus\{s\})};(s\vee(T\setminus\{s\}))_s) - x_{s\setminus r}b(r^{T\setminus\{s\}};(T\setminus\{s\})_r))\\
= \quad & \sum_{s\in T}(-1)^{\lambda(s\setminus r;T\setminus r)+i}(y_{s\setminus r}b(s^{s\vee(T\setminus\{s\})};(s\vee(T\setminus\{s\}))_s) - x_{s\setminus r}b(r^{T\setminus\{s\}};(T\setminus\{s\})_r)).
\end{aligned}
$$

On the other hand since $\mathscr{L}$ is a distributive lattice, the interval $[r, r^T]$ is a Boolean lattice. Hence $q \in T_r$, $r^T \setminus q = s \setminus r$ and $r^T \setminus T_r = T \setminus r$. Furthermore, we have

$$(s\vee(T\setminus\{s\}))_s = T_r \setminus q$$

and

$$r^{T\setminus\{s\}} = q, \quad (T\setminus\{s\})_r = q \wedge (T_r \setminus \{q\}).$$

These facts together with Theorem 3.1 yields

$$
\begin{aligned}
&(5) \qquad \partial_i(\pi_i(\sigma(\widetilde{b}(r;T)))) = \partial_i((-1)^i b(r^T;T_r))\\
= \quad & \sum_{q\in T_r}(-1)^{\lambda(r^T\setminus q;T_r\setminus r_T)+i}(y_{r^T\setminus q}b(r^T;T_r\setminus q) - x_{r^T\setminus q}b(q;q\wedge(T_r\setminus\{q\}))\\
= \quad & \sum_{s\in T}(-1)^{\lambda(s\setminus r;T\setminus r)+i}(y_{s\setminus r}b(s^{s\vee(T\setminus\{s\})};(s\vee(T\setminus\{s\}))_s) - x_{s\setminus r}b(r^{T\setminus\{s\}};(T\setminus\{s\})_r)).
\end{aligned}
$$

¿From (4) and (5) one sees that π is an isomorphism of complexes. $\square$

Let $\widetilde{\mathbb{T}}$ and $\widetilde{\mathbb{F}}$ be the minimal free resolutions of $H_{\widetilde{\mathscr{J}}}$ and $H_{\widetilde{\mathscr{L}}}$ as described in Theorem 3.1, and let $\iota : \widetilde{\mathbb{T}} \to \widetilde{\mathbb{F}}$ be the injective complex homomorphism which maps the basis elements $\widetilde{b}(r;T)$ of $\mathbb{T}$ to the corresponding basis elements of $\mathbb{F}$. Then we have the following sequence of complex homomorphisms:

$$\widetilde{\mathbb{T}} \xrightarrow{\iota} \widetilde{\mathbb{F}} \xrightarrow{\sigma} \sigma(\widetilde{\mathbb{F}}) \xrightarrow{\pi} \mathbb{F}.$$

Let ψ be the map from $\mathrm{Tor}(K, H_{\widetilde{\mathscr{J}}})$ to $\mathrm{Tor}(K, H_{\mathscr{L}})$ induced by $\pi\circ\sigma\circ\iota$.

As a consequence of the previous proposition, we now have:

Corollary 3.7. *For each $i \geq 0$ the map $\psi_i : \mathrm{Tor}_i(K, H_{\widetilde{\mathscr{J}}}) \to \mathrm{Tor}_i(K, H_{\mathscr{L}})$ is injective and maps the basis elements $\widetilde{b}(r;T)$ of $\mathrm{Tor}_i(K, H_{\widetilde{\mathscr{J}}})$ to the basis elements $(-1)^{|T|}b(r^T;T_r)$ of $\mathrm{Tor}_i(K, H_{\mathscr{L}})$.*

Now we are ready to present one of the main results of this section:

Theorem 3.8. *Let $\mathscr{L}$ be a finite distributive lattice, $\mathscr{I}$ a poset ideal and $\mathscr{J}$ a poset coideal of $\mathscr{L}$ such that $\mathscr{I} \cup \mathscr{J} = \mathscr{L}$. Then $H_{\mathscr{I}\cap\mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$ if and only if for each pair $p, q \in \mathscr{L}$ with $q \in N(p)$, either $p \in \mathscr{I}$ or $q \in \mathscr{J}$.*

Proof. We may assume that $|\mathscr{L}| > 1$, because otherwise the assertions are trivial.

Notice that $H_{\mathscr{I}\cap\mathscr{J}} \subseteq H_{\mathscr{I}} \cap H_{\mathscr{J}}$ holds always, and $H_{\mathscr{I}} \cap H_{\mathscr{J}} \subseteq H_{\mathscr{I}\cap\mathscr{J}}$ if and only if all generators of $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ have degree r, where r is the rank of $\mathscr{L}$.

Consider the long exact Tor-sequence

$$\begin{array}{llll}\longrightarrow \operatorname{Tor}_1(K,H_{\mathscr{I}})\oplus \operatorname{Tor}_1(K,H_{\mathscr{J}}) & \xrightarrow{\beta_1} \operatorname{Tor}_1(K,H_{\mathscr{I}}+H_{\mathscr{J}}) & \xrightarrow{\alpha_1} \operatorname{Tor}_0(K,H_{\mathscr{I}}\cap H_{\mathscr{J}}) \\ \longrightarrow \operatorname{Tor}_0(K,H_{\mathscr{I}})\oplus \operatorname{Tor}_0(K,H_{\mathscr{J}}) & \longrightarrow \operatorname{Tor}_0(K,H_{\mathscr{I}}+H_{\mathscr{J}}) & \longrightarrow \quad 0\end{array}$$

arising from the short exact sequence

$$0 \longrightarrow H_{\mathscr{I}} \cap H_{\mathscr{J}} \longrightarrow H_{\mathscr{I}} \oplus H_{\mathscr{J}} \longrightarrow H_{\mathscr{I}} + H_{\mathscr{J}} \longrightarrow 0.$$

Since $\mathscr{L} = \mathscr{I} \cup \mathscr{J}$, it follows that $H_{\mathscr{L}} = H_{\mathscr{I}} + H_{\mathscr{J}}$, and since $H_{\mathscr{L}}$ has an r-linear resolution we have $\operatorname{Tor}_i(K,H_{\mathscr{L}})_{i+j} = 0$ if $j \neq r$, and $\operatorname{Tor}_1(K,H_{\mathscr{L}})_{1+r} \neq 0$. Thus we see that all generators of $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ have degree r if and only if α_1 is a zero map, i.e, β_1 is a surjective map.

By Lemma 3.3 we have that the K-vector space $\beta_1(\operatorname{Tor}_1(K,H_{\mathscr{I}}))$ is spanned by the elements $b(p;\{q\})$ with $p \in \mathscr{I}$ and $q \in N(p)$, and by Corollary 3.7 we have that the vector space $\beta_1(\operatorname{Tor}_1(K,H_{\mathscr{J}}))$ is spanned by the elements $b(q^{\{p\}};\{p\}_q) = b(p;\{q\})$ with $q \in \mathscr{J}$ and $p \in M(q)$. It follows that the image of β_1 is spanned by the subset

$$B' = \{b(p;\{q\})\colon p \in \mathscr{I} \text{ and } q \in N(p), \text{ or } q \in \mathscr{J} \text{ and } q \in N(p)\}$$

of the basis

$$B = \{b(p;\{q\})\colon p \in \mathscr{L} \text{ and } q \in N(p)\}$$

of $\operatorname{Tor}_1(K,H_{\mathscr{L}})$. Therefore, β_1 is surjective if and only if $B' = B$. This implies the assertion. □

It is clear that if $\mathscr{I} \cap \mathscr{J} \neq \emptyset$ and $H_{\mathscr{I}\cap\mathscr{J}} \neq H_{\mathscr{I}} \cap H_{\mathscr{J}}$, then not all generators of $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ have the same degree. Therefore in this case, the ideal $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has no linear resolution. However in case $\mathscr{I} \cap \mathscr{J} \neq \emptyset$ and $H_{\mathscr{I}\cap\mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$ we have

Theorem 3.9. *Let $\mathscr{L}$ be a finite distributive lattice, $\mathscr{I}$ a poset ideal and $\mathscr{J}$ a poset coideal of $\mathscr{L}$ such that $\mathscr{I} \cup \mathscr{J} = \mathscr{L}$. If $H_{\mathscr{I}\cap\mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$, then the following statements are equivalent:*

1. *$H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has a linear resolution;*
2. *for each element $p \in \mathscr{L}$, either $p \in \mathscr{I}$ or $\wedge N(p) \in \mathscr{J}$;*
3. *for each element $r \in \mathscr{L}$, either $r \in \mathscr{J}$ or $\vee M(r) \in \mathscr{I}$.*

Proof. We may assume that $|\mathscr{L}| > 1$.

(2) ⇒ (3): Assume there exists some element $r \in \mathscr{L}$ such that $r \notin \mathscr{J}$ and $p = \vee M(r)$ does not belong to $\mathscr{I}$. Since $\mathscr{L}$ is a distributive lattice, the interval $[r,p]$ is a Boolean lattice. Hence $\wedge N(p) = r$. Therefore we have $p \notin \mathscr{I}$ and $r = \wedge N(p) \notin \mathscr{J}$, a contradiction.

By the same argument, one sees that (3) implies (2).

Now, we prove that the conditions (1) and (2) are equivalent. Consider the long exact Tor-sequence

$$\begin{aligned} \cdots \to \operatorname{Tor}_{i+1}(K,H_{\mathscr{I}})\oplus \operatorname{Tor}_{i+1}(K,H_{\mathscr{J}}) &\xrightarrow{\beta_{i+1}} \operatorname{Tor}_{i+1}(K,H_{\mathscr{L}}) \xrightarrow{\alpha_{i+1}} \operatorname{Tor}_{i}(K,H_{\mathscr{I}}\cap H_{\mathscr{J}}) \\ \longrightarrow \operatorname{Tor}_{i}(K,H_{\mathscr{I}})\oplus \operatorname{Tor}_{i}(K,H_{\mathscr{J}}) &\longrightarrow \operatorname{Tor}_{i}(K,H_{\mathscr{L}}) \longrightarrow \cdots \end{aligned}$$

arising from the short exact sequence

$$0 \longrightarrow H_{\mathscr{I}}\cap H_{\mathscr{J}} \longrightarrow H_{\mathscr{I}}\oplus H_{\mathscr{J}} \longrightarrow H_{\mathscr{L}} \longrightarrow 0.$$

Here we used that $\mathscr{I}\cup\mathscr{J}=\mathscr{L}$, so that $H_{\mathscr{I}}+H_{\mathscr{J}}=H_{\mathscr{L}}$. Let $r=\operatorname{rank}\mathscr{L}$. Since the ideal $H_{\mathscr{I}}\cap H_{\mathscr{J}}=H_{\mathscr{I}\cap\mathscr{J}}$ is generated in degree r, it has a linear resolution if and only if

$$\operatorname{Tor}_{i+1}(K,H_{\mathscr{L}})_j \xrightarrow{\alpha_{i+1,j}} \operatorname{Tor}_{i}(K,H_{\mathscr{I}}\cap H_{\mathscr{J}})_j$$

is the zero map for all $j\neq i+r$ and all $i\geq 0$, since the ideals $H_{\mathscr{I}}$ and $H_{\mathscr{J}}$ have r-linear resolutions. Since the ideal $H_{\mathscr{L}}$ has an r-linear resolution, the map $\alpha_{i+1,j}=0$ for $j\neq i+1+r$. Hence $H_{\mathscr{I}}\cap H_{\mathscr{J}}$ has a linear resolution if and only if $\alpha_{i+1,i+1+r}=0$ for all $i\geq 0$, and this is the case if and only if $\beta_{i+1,i+1+r}$ is surjective for all $i\geq 0$.

We argue as in the proof of Theorem 3.8. The set

$$B_{i+1}=\{b(p;S)\colon\, p\in\mathscr{L} \text{ and } S\subset N(p),\, |S|=i+1\}$$

is a K-basis of $\operatorname{Tor}_{i+1}(K,H_{\mathscr{L}})_{i+1+r}$. Using Lemma 3.3 and Corollary 3.7 we see that the set

$$\begin{aligned} B'_{i+1} &= \{b(p;S)\colon\, p\in\mathscr{I} \text{ and } S\subset N(p),\, |S|=i+1\} \\ &\cup \{b(r^T;T_r)\colon\, r\in\mathscr{J} \text{ and } T\subset M(r),\, |T|=i+1\} \end{aligned}$$

spans the image of $\beta_{i+1,i+1+r}$. Thus $\beta_{i+1,i+1+r}$ is surjective if and only if $B'_{i+1}=B_{i+1}$ for all $i>0$. Note that $B'_{i+1}\subset B_{i+1}$. Suppose condition (2) holds, and let $b(p;S)\in B_{i+1}$. If $p\in\mathscr{I}$, then $b(p;S)\in B'_{i+1}$. If $p\notin\mathscr{I}$, then $p\in\mathscr{J}$. Let $r=\wedge S$. It follows from condition (2) that $r\in\mathscr{J}$. Let T be the set of upper neighbors of r in the interval $[r,p]$. Then $p=r^T$ and $S=T_r$, and hence $b(p;S)=b(r^T;T_r)$ belongs to B'_{i+1}.

Conversely assume that $B'_{i+1}=B_{i+1}$. In particular, for all $p\in\mathscr{L}$ we have $b(p;N(p))\in B'_{|N(p)|}$. So either $p\in\mathscr{I}$ or there is some $r\in\mathscr{J}$ and $T\subset M(r)$ such that $p=r^T$ and $T_r=N(p)$. Since $\wedge T_r=r$, it follows that $\wedge N(p)=r$ which is in $\mathscr{J}$. $\square$

Up to now, we always assume that $\mathscr{I}\cup\mathscr{J}=\mathscr{L}$ and $\mathscr{I}\cap\mathscr{J}\neq\emptyset$. Now we consider the case $\mathscr{I}\cup\mathscr{J}=\mathscr{L}$ and $\mathscr{I}\cap\mathscr{J}=\emptyset$. In this case, $H_{\mathscr{I}\cap\mathscr{J}}\neq H_{\mathscr{I}}\cap H_{\mathscr{J}}$. However, we have:

Theorem 3.10. *Let $\mathscr{L}$ be a finite distributive lattice, $\mathscr{I}$ a poset ideal and $\mathscr{J}$ a poset coideal of $\mathscr{L}$. If $\mathscr{I}\cup\mathscr{J}=\mathscr{L}$ and $\mathscr{I}\cap\mathscr{J}=\emptyset$, then the ideal $H_{\mathscr{I}}\cap H_{\mathscr{J}}$ has a linear resolution.*

This theorem follows immediately from the following two lemmata.

Lemma 3.11. *Under the assumption of Proposition 3.10, the ideal $H_{\mathscr{I}}\cap H_{\mathscr{J}}$ is generated by the monomials* $\operatorname{lcm}(u_p,u_q)$ *with $p\in\mathscr{J}$, $q\in\mathscr{I}$ and $q\in N(p)$. In particular, all generators of $H_{\mathscr{I}}\cap H_{\mathscr{J}}$ are of degree* $\operatorname{rank}\mathscr{L}+1$.

Proof. Since $\mathscr{I} \cap \mathscr{J} = \emptyset$, all generators of $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ have degree greater than the rank of $\mathscr{L}$. Let $H = \langle \operatorname{lcm}(u_p, u_q) : p \in \mathscr{J}, q \in \mathscr{I} \text{ and } q \in N(p) \rangle$. It is clear that $H \subseteq H_{\mathscr{I}} \cap H_{\mathscr{J}}$. Since $H_{\mathscr{I}}$ and $H_{\mathscr{J}}$ both are monomial ideals, the intersection $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ is again a monomial ideal. Let m be any monomial in $H_{\mathscr{I}} \cap H_{\mathscr{J}}$. Then there exist $r \in \mathscr{I}$ and $s \in \mathscr{J}$, such that $u_r | m$ and $u_s | m$. Let C be any chain between $r \wedge s$ and $r \vee s$. Since $\mathscr{I}$ is a poset ideal and $\mathscr{J}$ is a poset coideal, and $r \in \mathscr{I}$, $s \in \mathscr{J}$, we have $r \wedge s \in \mathscr{I}$ and $r \vee s \in \mathscr{J}$. Hence there exist $p, q \in C$ such that $q \in \mathscr{I}$, $p \in \mathscr{J}$ and q is a lower neighbor of p. We claim $\operatorname{lcm}(u_p, u_q) | m$. To see this, we write $m = \prod_{i=1}^{n} x^{a_i} y^{b_i}$ as $m_x m_y$ where $m_x = \prod_{i=1}^{n} x^{a_i}$ and $m_y = \prod_{i=1}^{n} y^{b_i}$, and as before we write $u_t = x_{\ell(t)} y_{P \setminus \ell(t)}$, where P is the set of join irreducible elements of $\mathscr{L}$. Since $\ell(p) \subseteq \ell(r \vee s)$, we have $x_{\ell(p)} | x_{\ell(r \vee s)}$. Since $u_r | m$ and $u_s | m$, we have $x_{\ell(r \vee s)} | m$ and hence $x_{\ell(p)} | m_x$. By the same argument we see that $y_{P \setminus \ell(q)} | m_y$. Hence $\operatorname{lcm}(u_p, u_q) = x_{\ell(p)} y_{P \setminus \ell(q)}$ divides m. $\square$

Lemma 3.12. *Let R be a polynomial ring over a field K, I and J ideals in R. Suppose I, J and $I + J$ have d-linear resolutions. If all elements of $G(I \cap J)$ have degree $d + 1$, then the ideal $I \cap J$ has a $(d+1)$-linear resolution.*

Proof. Consider the long exact Tor-sequence

$$\begin{aligned} \cdots \to \operatorname{Tor}_{i+1}(K, I) \oplus \operatorname{Tor}_{i+1}(K, J) &\xrightarrow{\beta_{i+1}} \operatorname{Tor}_{i+1}(K, I+J) \xrightarrow{\alpha_{i+1}} \operatorname{Tor}_i(K, I \cap J) \\ \longrightarrow \operatorname{Tor}_i(K, I) \oplus \operatorname{Tor}_i(K, J) &\longrightarrow \operatorname{Tor}_i(K, I+J) \longrightarrow \cdots. \end{aligned}$$

Since I, J and $I + J$ have d-linear resolutions. It follows that $\operatorname{Tor}_i(K, I)_j = \operatorname{Tor}_i(K, J)_j = 0$ for any $j \neq i + d$ and $\operatorname{Tor}_{i+1}(K, I+J)_j = 0$ for any $j \neq i + 1 + d$. Hence $\operatorname{Tor}_i(K, I \cap J)_j = 0$ for $j < i + d$ or $j > i + 1 + d$. Since $I \cap J$ is generated in degree $d + 1$, we have $\operatorname{Tor}_i(K, I \cap J)_j = 0$ for $j = i + d$. Therefore $\operatorname{Tor}_i(K, I \cap J)_j = 0$ for any $j \neq i + 1 + d$. Hence $I \cap J$ has a $(d+1)$-linear resolution. $\square$

In the remaining of this section we discuss some special classes of segments of a finite distributive lattice $\mathscr{L}$ to which our results apply and where some additional information can be obtained . As we have already seen, for any segment $\mathscr{S}$ there exist a poset ideal $\mathscr{I}$ and a poset coideal $\mathscr{J}$ such that $\mathscr{S} = \mathscr{I} \cap \mathscr{J}$. Now Let $\mathscr{L}$ be a finite distributive lattice of rank r. We consider a special class of segments $\mathscr{S}$ of $\mathscr{L}$ which consisting of all elements p in $\mathscr{L}$ such that $i \leq \operatorname{rank} p \leq j$ for some i and j with $0 \leq i \leq j \leq r$. We denote it by $\mathscr{L}_{i,j}$.

Lemma 3.13. *Let $\mathscr{L}$ be a finite distributive lattice of rank r, and let $\mathscr{L}_{i,j}$ be a segment of $\mathscr{L}$. Then there exists a poset ideal $\mathscr{I}$ and a poset coideal $\mathscr{J}$ of $\mathscr{L}$ such that $H_{\mathscr{L}_{i,j}} = H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$.*

Proof. Let $\mathscr{I} = \{p \in \mathscr{L} : \operatorname{rank} p \leq j\}$, and $\mathscr{J} = \{p \in \mathscr{L} : \operatorname{rank} p \geq i\}$. Then $\mathscr{I}$ is a poset ideal, $\mathscr{J}$ is a poset coideal of $\mathscr{L}$, and $\mathscr{L}_{i,j} = \mathscr{I} \cap \mathscr{J}$. It remains to show that $H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$. Let $p, q \in \mathscr{L}$ and $q \in N(p)$. If $p \notin \mathscr{I}$, then $\operatorname{rank} p > j$. Hence $\operatorname{rank} q = \operatorname{rank} p - 1 \geq j \geq i$, i.e., $q \in \mathscr{J}$. The assertion follows from Proposition 3.8. $\square$

With the assumptions and notation of the previous lemma, the ideal $H_{\mathscr{L}_{i,j}}$ has a linear resolution if and only if $H_{\mathscr{I}} \cap H_{\mathscr{J}}$ has a linear resolution.

Corollary 3.14. *Let $\mathscr{L} \neq \{\hat{0}, \hat{1}\}$ be a finite distributive lattice and $\mathscr{S} = \mathscr{L} \setminus \{\hat{0}, \hat{1}\}$. Then*

1. *$H_{\mathscr{S}}$ has a linear resolution if and only if $\mathscr{L}$ is not a Boolean lattice;*
2. *in case $\mathscr{L}$ is the Boolean lattice $\mathscr{B}_r$, the ideal $H_{\mathscr{S}}$ has the following minimal free resolution:*

$$\mathbb{T}\colon 0 \longrightarrow T_{r-1} \longrightarrow \cdots T_1 \longrightarrow T_0 \longrightarrow H_{\mathscr{S}} \longrightarrow 0.$$

with $T_i = S^{\binom{r}{i}(2^{r-i}-2)}(-r-i)$ for $i = 0, \ldots, r-2$ and $T_{r-1} = S(-2r)$.

Proof. (1) Let $\mathscr{I} = \{p \in \mathscr{L} : \operatorname{rank} p \leq \operatorname{rank} \mathscr{L} - 1\}$, and $\mathscr{J} = \{p \in \mathscr{L} : \operatorname{rank} p \geq 1\}$. Then by Lemma 3.13, $\mathscr{I}$ and $\mathscr{J}$ are the poset ideal and poset coideal of $\mathscr{L}$ such that $H_{\mathscr{S}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$. The distributive lattice $\mathscr{L}$ is a Boolean lattice if and only if the meet of all lower neighbors of $\hat{1}$ is $\hat{0}$. Since $\hat{1}$ is the only element which is not in $\mathscr{I}$, and $\hat{0}$ is the only element which is not in $\mathscr{J}$, by using the Theorem 3.9, we have $H_{\mathscr{S}}$ has a linear resolution if and only if $\mathscr{L}$ is not a Boolean lattice.

(2) Choose $\mathscr{I}$ and $\mathscr{J}$ as in the proof of (1). Hence $H_{\mathscr{S}} = H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$. We consider long exact Tor-sequence as in the proof of Theorem 3.9. Notice that $(\hat{1}, N(\hat{1}))$ is the only pair with the form (p, S) with $p \in \mathscr{B}_r$ and $S \subset N(p)$ such that $p \notin \mathscr{I}$ and $\wedge S \notin \mathscr{J}$. It follows that the map β_i is surjective for $i < r$. Hence for all $i < r-1$ we have the exact sequence

$$(6) \quad 0 \longrightarrow \operatorname{Tor}_i(K, H_{\mathscr{S}}) \longrightarrow \operatorname{Tor}_i(K, H_{\mathscr{I}}) \oplus \operatorname{Tor}_i(K, H_{\mathscr{J}}) \longrightarrow \operatorname{Tor}_i(K, H_{\mathscr{B}_r}) \longrightarrow 0,$$

and so

$$b_i(H_{\mathscr{S}}) = b_i(H_{\mathscr{I}}) + b_i(H_{\mathscr{J}}) - b_i(H_{\mathscr{B}_r}) \quad \text{for} \quad i < r-1,$$

where $b_i(I)$ is the i-th Betti number of the ideal I. By using Theorem 3.1, Corollary 3.7 and the combinatorial fact that each Boolean lattice $\mathscr{B}_r$ contains $\binom{r}{i}2^{r-i}$ Boolean sublattices $\mathscr{B}_i$, we have $b_i(H_{\mathscr{I}}) = b_i(H_{\mathscr{J}}) = \binom{r}{i}2^{r-i} - \binom{r}{i}$ and $b_i(H_{\mathscr{L}}) = \binom{r}{i}2^{r-i}$. Hence

$$b_i(H_{\mathscr{S}}) = 2(\binom{r}{i}2^{r-i} - \binom{r}{i}) - \binom{r}{i}2^{r-i} = \binom{r}{i}(2^{r-i}-2),$$

for any $i < r-1$. It also follows from (6) that the resolution of $H_{\mathscr{S}}$ is linear up to homological degree $r-2$.

Now let $i = r-1$. Since $\operatorname{Tor}_r(K, H_{\mathscr{I}}) = \operatorname{Tor}_r(K, H_{\mathscr{J}}) = 0$, we get the exact sequence

$$\begin{aligned} 0 \to \operatorname{Tor}_r(K, H_{\mathscr{B}_r}) \to \quad & \operatorname{Tor}_{r-1}(K, H_{\mathscr{S}}) \quad \to \operatorname{Tor}_{r-1}(K, H_{\mathscr{I}}) \oplus \operatorname{Tor}_{r-1}(K, H_{\mathscr{J}}) \\ \longrightarrow \quad & \quad \operatorname{Tor}_{r-1}(K, H_{\mathscr{B}_r}) \to 0. \end{aligned}$$

Since $\dim_K \operatorname{Tor}_{r-1}(K, H_{\mathscr{I}}) \oplus \operatorname{Tor}_{r-1}(K, H_{\mathscr{J}}) = \dim_K \operatorname{Tor}_{r-1}(K, H_{\mathscr{J}}) = 2r$, it follows that $\operatorname{Tor}_{r-1}(K, H_{\mathscr{I}}) \oplus \operatorname{Tor}_{r-1}(K, H_{\mathscr{J}}) \to \operatorname{Tor}_{r-1}(K, H_{\mathscr{B}_r})$ is an isomorphism. Hence

$$\operatorname{Tor}_{r-1}(K, H_{\mathscr{S}}) \cong \operatorname{Tor}_r(K, H_{\mathscr{B}_r}) \cong K(-2r),$$

as desired. □

Using Lemma 3.13 and Theorem3.9, we have the following two facts:

Corollary 3.15. *Let $\mathscr{L}$ be a finite distributive lattice, and i an integer. If $|\mathscr{L}_{i,i}| > 1$, then the ideal $H_{\mathscr{L}_{i,i}}$ has no linear resolution.*

Proof. Let $\mathscr{I} = \{p \in \mathscr{L} : \operatorname{rank} p \leq i\}$ and $\mathscr{J} = \{p \in \mathscr{L} : \operatorname{rank} p \geq i\}$. Then by Lemma 3.13 we have $H_{\mathscr{L}_{i,i}} = H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$. Since $\mathscr{L}$ is distributive, there exist elements u and v in $\mathscr{L}_{i,i}$ such that $\operatorname{rank}(u \vee v) = i + 1$. Hence $\operatorname{rank}(\wedge N(u \vee v)) < \operatorname{rank} u = i$. Therefore we have $u \vee v \notin \mathscr{I}$ and $\wedge N(u \vee v) \notin \mathscr{J}$. By Theorem 3.9, $H_{\mathscr{L}_{i,i}}$ has no linear resolution. □

Corollary 3.16. *If $\mathscr{L}$ is a finite planar distributive lattice of rank r, then the ideal $H_{\mathscr{L}_{i,j}}$ has a linear resolution, if $i < j$.*

Proof. Let $\mathscr{I} = \{p \in \mathscr{L} : \operatorname{rank} p \leq j\}$ and $\mathscr{J} = \{p \in \mathscr{L} : \operatorname{rank} p \geq i\}$. Then $H_{\mathscr{L}_{i,j}} = H_{\mathscr{I} \cap \mathscr{J}} = H_{\mathscr{I}} \cap H_{\mathscr{J}}$. Since $\mathscr{L}$ is a planar distributive lattice, each element p in $\mathscr{L}$ has at most two lower neighbors. Hence $\operatorname{rank} \wedge N(p) \geq p - 2$. Thus if $p \notin \mathscr{I}$, then $\operatorname{rank} p > j$. Therefore since $i < j$, we have $\operatorname{rank} q \geq \operatorname{rank} p - 2 \geq i$, i.e., $q \in \mathscr{J}$. By Theorem 3.9, $H_{\mathscr{L}_{i,j}}$ has a linear resolution. □

REFERENCES

[1] D: Bayer and B. Sturmfels, Cellular resolutions of monomial modules, *J. reine angew. Math.* **502** (1998), 123–140.
[2] W. Bruns and J. Herzog, "Cohen–Macaulay rings," Revised Edition, Cambridge University Press, 1996.
[3] J. Eagon and V. Reiner, Resolutions of Stanley–Reisner rings and Alexander duality, *J. Pure Appl. Algebra* **130** (1998), 265–275.
[4] S. Faridi, The facet ideal of a simplicial complex, Manuscripta Mathematica 109, (2002), 159-174.
[5] T. Hibi, "Algebraic Combinatorics on Convex Polytopes," Carslew Publications, 1992.
[6] J. Herzog and T. Hibi, Distributive Lattices, Bipartite Graphs and Alexander Duality, preprint, 2003.
[7] J. Herzog, T. Hibi and X. Zheng, Dirac's theorem on chordal graphs and Alexander duality, to appear in YEUJC.
[8] J. Herzog, T. Hibi and X. Zheng, The monomial ideal of a finite meet-semilattice, preprint, 2003.
[9] R. P. Stanley, "Enumerative Combinatorics, Volume I," Wadsworth & Brooks/Cole, Monterey, CA, 1986.
[10] R. P. Stanley, "Combinatorics and Commutative Algebra," Second Edition, Birkhäuser, Boston, MA, 1996.
[11] X. Zheng, Resolutions of facet ideals, to appear in Communication in Alg.

XINXIAN ZHENG, FACHBEREICH MATHEMATIK UND INFORMATIK, UNIVERSITÄT DUISBURG-ESSEN, 45117 ESSEN, GERMANY
E-mail address: `xinxian.zheng@uni-essen.de`